



## Forcing versus feedback: Epidemic malaria and monsoon rains in northwest India

**Author(s):** Laneri K, Bhadra A, Ionides EL, Bouma M, Dhiman RC, Yadav RS, Pascual M  
**Year:** 2010  
**Journal:** PLoS Computational Biology. 6 (9): e1000898

### Abstract:

Malaria epidemics in regions with seasonal windows of transmission can vary greatly in size from year to year. A central question has been whether these interannual cycles are driven by climate, are instead generated by the intrinsic dynamics of the disease, or result from the resonance of these two mechanisms. This corresponds to the more general inverse problem of identifying the respective roles of external forcings vs. internal feedbacks from time series for nonlinear and noisy systems. We propose here a quantitative approach to formally compare rival hypotheses on climate vs. disease dynamics, or external forcings vs. internal feedbacks, that combines dynamical models with recently developed, computational inference methods. The interannual patterns of epidemic malaria are investigated here for desert regions of northwest India, with extensive epidemiological records for *Plasmodium falciparum* malaria for the past two decades. We formulate a dynamical model of malaria transmission that explicitly incorporates rainfall, and we rely on recent advances on parameter estimation for nonlinear and stochastic dynamical systems based on sequential Monte Carlo methods. Results show a significant effect of rainfall in the inter-annual variability of epidemic malaria that involves a threshold in the disease response. The model exhibits high prediction skill for yearly cases in the malaria transmission season following the monsoonal rains. Consideration of a more complex model with clinical immunity demonstrates the robustness of the findings and suggests a role of infected individuals that lack clinical symptoms as a reservoir for transmission. Our results indicate that the nonlinear dynamics of the disease itself play a role at the seasonal, but not the interannual, time scales. They illustrate the feasibility of forecasting malaria epidemics in desert and semi-arid regions of India based on climate variability. This approach should be applicable to malaria in other locations, to other infectious diseases, and to other nonlinear systems under forcing.

**Source:** <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2932675>

### Resource Description

#### Early Warning System:

resource focus on systems used to warn populations of high temperatures, extreme weather, or other elements of climate change to prevent harm to health

A focus of content

#### Exposure :

weather or climate related pathway by which climate change affects health

# Climate Change and Human Health Literature Portal

Extreme Weather Event, Precipitation

## **Geographic Feature:**

resource focuses on specific type of geography

None or Unspecified

## **Geographic Location:**

resource focuses on specific location

Non-United States

**Non-United States:** Asia

**Asian Region/Country:** India

## **Health Impact:**

specification of health effect or disease related to climate change exposure

Infectious Disease

**Infectious Disease:** Vectorborne Disease

**Vectorborne Disease:** Mosquito-borne Disease

**Mosquito-borne Disease:** Malaria

## **Mitigation/Adaptation:**

mitigation or adaptation strategy is a focus of resource

Adaptation

## **Model/Methodology:**

type of model used or methodology development is a focus of resource

Exposure Change Prediction, Outcome Change Prediction

**Population of Concern:** A focus of content

## **Population of Concern:**

populations at particular risk or vulnerability to climate change impacts

Children

## **Resource Type:**

format or standard characteristic of resource

Research Article

## **Timescale:**

time period studied

Time Scale Unspecified

## **Vulnerability/Impact Assessment:**

resource focus on process of identifying, quantifying, and prioritizing vulnerabilities in a system

A focus of content